

POLARIZATIONS OF PRYM VARIETIES OF PAIRS OF COVERINGS

H. LANGE AND S. RECILLAS

Sevin Recillas died on June 20, 2005.

ABSTRACT. To any pair of coverings $f_i : X \rightarrow X_i, i = 1, 2$ of smooth projective curves one can associate an abelian subvariety of the Jacobian J_X , the Prym variety $P(f_1, f_2)$ of the pair (f_1, f_2) . In some cases we can compute the type of the restriction of the canonical principal polarization of J_X . We obtain 2 families of Prym-Tyurin varieties of exponent 6.

1. INTRODUCTION

Let $f : X \rightarrow Y$ be a finite morphism of smooth projective curves. The complement of the abelian subvariety f^*J_Y in the canonically polarized Jacobian J_X of X is called the Prym variety $P(f)$ of f . In [3] we introduced an analogous notion for two morphisms of curves. To be more precise, suppose that we are given a commutative diagram of finite morphisms of smooth projective curves:

$$\begin{array}{ccc} & X & \\ f_1 \swarrow & & \searrow f_2 \\ X_1 & & X_2 \\ g_1 \searrow & & \swarrow g_2 \\ & Y & \end{array} \quad (1.1)$$

such that g_1 and g_2 do not both factorize via the same morphism $Y' \rightarrow Y$ of degree ≥ 2 . Then the *Prym variety* $P(f_1, f_2)$ of the pair (f_1, f_2) is defined to be the complement of the abelian variety $f_2^*(P(g_2))$ in $P(f_1)$ with respect to the canonical polarization.

By definition $P(f_1, f_2)$ is an abelian subvariety of the Jacobian J_X of the curve X . In general it is difficult to determine the type of the restriction of a polarization to an abelian subvariety. It is the aim of this note to determine the type of the restriction of the canonical polarization of J_X to $P(f_1, f_2)$ in some cases, namely for $Y = \mathbb{P}^1$ and $\deg f_1$ and $\deg f_2$ are prime to each other. As a special case we obtain 2 families of Prym-Tyurin varieties of exponent 6 in any dimension ≥ 4 respectively 5. Note that the correspondences defining $P(f_1, f_2)$

2000 *Mathematics Subject Classification*. Primary: 14K05; Secondary: 14H40.

Key words and phrases. Prym variety, Prym-Tyurin variety, polarization.

Supported by DAAD and Conacyt 40033-F.

are not fixed point free. So one cannot apply Kanev's theorem for the proof.

In the last section we show that the Abel-Prym map of these abelian varieties is injective apart from the fact that all ramification points of f_1 are mapped to one point. Hence we obtain irreducible curves of low class in $P(f_1, f_2)$. Moreover this implies that the Seshadri constant of $P(f_1, f_2)$ is small in these cases.

2. RESTRICTING POLARIZATIONS

Let (S, L) be a polarized abelian variety over an algebraically closed field of characteristic 0. As usual let $K(L)$ denote the kernel of the associated homomorphism $\phi_L : S \rightarrow \hat{S}$, $x \mapsto t_x^* L \otimes L^{-1}$. The polarization L is of type $(d_1, \dots, d_g)$ if and only if $K(L) \simeq (\mathbb{Z}/d_1\mathbb{Z} \times \dots \times \mathbb{Z}/d_g\mathbb{Z})^2$.

Now let A be an abelian subvariety of S with canonical embedding $\iota_A : A \rightarrow S$. Let

$$P := \ker(\hat{\iota}_A \circ \phi_L : S \rightarrow \hat{A})^0 \quad (2.1)$$

denote the complementary abelian subvariety of S . Let $L_A := L|_A$ and $L_P := L|_P$ denote the restriction of the polarization L to A and P . If L is a principal polarization, it is well known that

$$K(L_A) \simeq K(L_P) \simeq A \cap P \quad (2.2)$$

from which the type of L_A and L_P can be deduced in many cases. For arbitrary polarizations the situation is more complicated. In this section we recall two generalizations of (2.2) which will be applied later and for the proof of which we refer to [2].

Proposition 2.1. $|K(L_P)| \cdot |K(L_A)| = |A \cap P|^2 \cdot |K(L)|$.

Proposition 2.2. *There is an exact sequence*

$$0 \rightarrow K(L) \cap P \rightarrow K(L_P) \rightarrow A \cap P \rightarrow 0.$$

In particular $|K(L_P)| = |K(L) \cap P| \cdot |A \cap P|$ and similarly for L_A .

3. RESTRICTION OF THE CANONICAL POLARIZATION TO $P(f_1, f_2)$

Let $f : X \rightarrow Y$ be a morphism of degree n of smooth projective curves over an algebraically closed field k . Denote by $J_X := \text{Pic}^0(X)$ and $J_Y := \text{Pic}^0(Y)$ the Jacobians of X and Y . Pulling back line bundles defines a homomorphism

$$f^* : J_Y \rightarrow J_X.$$

f^* has finite kernel and is an embedding if and only if f does not factor via a cyclic étale cover of degree ≥ 2 (see [1], Proposition 11.4.3). The norm map of line bundles defines a homomorphism

$$N_f : J_X \longrightarrow J_Y.$$

The *Prym variety* $P(f)$ of the morphism f is defined to be the abelian subvariety

$$P(f) := \ker(N_f)^0$$

of J_X where the 0 means the connected component containing 0.

Now suppose that we are given a commutative diagram of finite morphisms of smooth projective curves:

$$\begin{array}{ccc} & X & \\ f_1 \swarrow & & \searrow f_2 \\ X_1 & & X_2 \\ g_1 \searrow & & \swarrow g_2 \\ & \mathbb{P}^1 & \end{array} \quad (3.1)$$

with the following properties:

- (a) g_1 and g_2 do not both factorize via the same morphism $Y' \rightarrow \mathbb{P}^1$ of degree ≥ 2 ,
- (b) X is the normalisation of the cartesian product $X_1 \times_{\mathbb{P}^1} X_2$.
- (c) $d_1 := \deg g_1$ and $d_2 := \deg g_2$ are prime to each other.

Then we have for the Prym variety $P = P(f_1, f_2)$, defined as the complement of $f_2^*(P(g_2))$ in $P(f_1)$ with respect to the restriction of the canonical polarization of J_X to $P(f_1)$ (see [3]):

Proposition 3.1. *Let Θ be the canonical principal polarization of J_X and denote*

$$L := \Theta|_{P(f_1)}, \quad L_{f_2^*P(g_2)} := L|_{f_2^*P(g_2)} = \Theta|_{f_2^*P(g_2)}, \quad L_P := L|_P = \Theta|_P.$$

Then

$$K(L_P) \simeq K(L) \oplus K(L_{f_2^*P(g_2)}).$$

Proof. $P(f_1)$ is by definition the complementary abelian subvariety of $f_1^*J_{X_1}$ in J_X with respect to Θ . According to [1], Proposition 11.4.3 $\Theta|_{f_1^*J_{X_1}}$ is of type $(d_2^{(1)}, \dots, d_2^{(s)}, d_2, \dots, d_2)$ where $d_2^{(i)}$ divides d_2 for $i = 1, \dots, s$ and $0 \leq s < \dim J_{X_1}$. Hence, according to (2.2) the polarization L is of type $(1, \dots, 1, d_2^{(1)}, \dots, d_2^{(s)}, d_2, \dots, d_2)$. In particular $K(L)$ is of exponent d_2 .

On the other hand $P(g_2) = J_{X_2}$. Again by [1], Proposition 11.4.3, $L_{f_2^*P(g_2)} = \Theta|_{f_2^*P(g_2)}$ is of type $(d_1^{(1)}, \dots, d_1^{(t)}, d_1, \dots, d_1)$ where $d_1^{(i)}$ divides d_1 for $i = 1, \dots, t$ and $0 \leq t < \dim J_{X_2}$. In particular $L_{f_2^*P(g_2)}$ is of exponent d_1 .

Now P is by definition the complementary abelian subvariety of $f_2^*P(g_2)$

in $P(f_1)$ with respect to the polarization L . Hence according to Proposition 2.2 there is an exact sequence

$$0 \longrightarrow K(L) \cap f_2^*P(g_2) \longrightarrow K(L_{f_2^*P(g_2)}) \longrightarrow P \cap f_2^*P(g_2) \longrightarrow 0$$

The assumption $\gcd(d_1, d_2) = 1$ implies by the above reasoning that $\gcd(|K(L)|, |K(L_{f_2^*P(g_2)})|) = 1$. Hence $K(L) \cap f_2^*P(g_2) = 0$ and thus

$$K(L_{f_2^*P(g_2)}) \simeq f_2^*P(g_2) \cap P.$$

But then Proposition 2.1 yields

$$|K(L_P)| = |K(L)| \cdot |K(L_{f_2^*P(g_2)})|$$

which implies the assertion, since d_1 and d_2 are prime to each other. $\square$

By symmetry reasons we also get with the same notation

Remark 3.2. *Denote*

$$M := \Theta|P(f_2), \quad M_{f_1^*P(g_1)} := M|f_1^*P(g_1), \quad M_P = \Theta|P = L_P.$$

Then

$$K(L_P) = K(M_P) \simeq K(M) \oplus K(M_{f_1^*P(g_1)}).$$

Recall that P is a Prym-Tyurin variety of exponent q in the Jacobian J_X if the restricted polarization $L_P = \Theta|P$ is the q -fold of a principal polarization of P . This is the case if and only if L_P is of type $(q, \dots, q)$.

Proposition 3.3. *Let the notation be as above. In particular we assume $\gcd(d_1, d_2) = 1$. If we assume moreover that f_1 and f_2 do not factorize via a cyclic unramified covering, then the following conditions are equivalent*

- (a) $P = P(f_1, f_2)$ is a Prym-Tyurin variety of exponent d_1d_2 in J_X .
- (b) $\dim(P) = g(X_1) = g(X_2)$.

Proof. Since f_1 does not factorize via an unramified covering L is of type $(1, \dots, 1, d_2, \dots, d_2)$ with $g(X_1)$ entries d_2 . Since f_2 does not factorize via an unramified covering, $L_{f_2^*P(g_2)}$ is of type $(d_1, \dots, d_1)$ with $g(X_2)$ entries d_1 . Hence according to Proposition 3.1 L_P is of type $(1, \dots, 1, d_1d_2, \dots, d_1d_2)$ if $g(X_1) = g(X_2)$ with $g(X_1)$ entries d_1d_2 and $\dim P - g(X_1)$ entries 1. In particular we have $\dim P \geq g(X_1)$. If $\dim P = g(X_1) = g(X_2)$, there is no entry 1 implying that (P, L_P) is a Prym-Tyurin variety of exponent d_1d_2 in J_X . Conversely, if one of the equations in (b) is not satisfied, the above reasoning implies that (P, L_P) cannot be a Prym-Tyurin variety in J_X . $\square$

We also need the following variation of Proposition 3.3:

Proposition 3.4. *Let the notation be as in Proposition 3.3 and assume now that as above f_1 does not factorize via an unramified covering, but f_2 is étale cyclic. Then the following conditions are equivalent*

- (a) $P = P(f_1, f_2)$ is a Prym-Tyurin variety of exponent d_1d_2 in J_X .

(b) $\dim(P) = g(X_1) = g(X_2) - 1$ and $d_1 = 3$.

Moreover in this case $d_1 = 3$ if $\dim P \geq 2$.

Proof. As above L is of type $(1, \dots, 1, d_2, \dots, d_2)$ with $g(X_1)$ entries d_2 . Since f_2 is étale cyclic of degree d_1 , $L_{f_2^*P(g_2)}$ is of type $(1, d_1, \dots, d_1)$ with $g(X_2) - 1$ entries d_1 . Hence according to Proposition 3.1 L_P is of type $(1, \dots, 1, d_1d_2, \dots, d_1d_2)$ if $g(X_1) = g(X_2) - 1$ with $g(X_1)$ entries d_1d_2 and $\dim P - g(X_1)$ entries 1. As in the proof of Proposition 3.3 one concludes that if $\dim(P) = g(X_1) = g(X_2) - 1$, then P is a Prym-Tyurin variety.

Moreover, since f_2 is cyclic étale, we have $g(X) = d_1g(X_2) - d_1 + 1$ and thus

$$\begin{aligned} g(X_2) - 1 = \dim P &= g(X) - g(X_1) - g(X_2) \\ &= (d_1 - 2)g(X_2) - d_1 + 2, \end{aligned}$$

implying $d_1 = 3$ if $g(X_2) = \dim P + 1 \geq 3$. Again, if the equation in (b) are not satisfied, it is easy to see that (P, L_P) cannot be a Prym-Tyurin variety in J_X . $\square$

4. TWO FAMILIES OF PRYM-TYURIN VARIETIES

In this section we analyse the conditions of Propositions 3.3 and 3.4 to obtain two families of Prym varieties of pairs of morphisms which are Prym-Tyurin varieties. Suppose we are given a commutative diagram (3.1). For any finite morphism f of smooth projective curves let $\delta(f)$ denote the degree of the ramification divisor. This is always an even number. Hence we may denote

$$\delta(g_i) = 2r_i \quad \text{and} \quad \delta(f_i) = 2s_i \quad \text{for} \quad i = 1, 2.$$

Of course r_1, r_2, s_1 and s_2 are not unrelated. Then Hurwitz' formula gives

$$g(X_i) = r_i - d_i + 1 \quad \text{for} \quad i = 1, 2$$

and

$$g(X) = d_2(r_1 - d_1) + s_1 + 1 = d_1(r_2 - d_2) + s_2 + 1.$$

This implies

$$\begin{aligned} \dim P &= d_1r_2 - d_1d_2 + d_1 + d_2 - r_1 - r_2 + s_2 - 1 \\ &= d_2r_1 - d_1d_2 + d_1 + d_2 - r_1 - r_2 + s_1 - 1 \end{aligned}$$

In particular we have

$$s_2 = d_2r_1 - d_1r_2 + s_1. \tag{4.1}$$

Theorem 4.1. *Let the assumptions be as in Proposition 3.3. In particular we assume that $\gcd(d_1, d_2) = 1$ and that f_1 and f_2 do not factorize via a cyclic étale covering. Without loss of generality we may assume $d_1 > d_2$.*

Then $P = P(f_1, f_2)$ is a Prym-Tyurin variety of dimension ≥ 4 in J_X if and only if

$$d_1 = 3, d_2 = 2, r_1 \geq 6, s_1 = r_2 = r_1 - 1 \text{ and } s_2 = 2.$$

In this case P is a Prym-Tyurin variety of dimension $r_1 - 2$ in J_X .

Proof. According to Proposition 3.3 P is a Prym-Tyurin variety of exponent $d_1 d_2$ in J_X if and only if $\dim P = g(X_1) = g(X_2)$ which is equivalent to

$$\begin{cases} r_1 - d_1 &= r_2 - d_2 \\ d_1 r_2 - d_1 d_2 + d_1 + d_2 - r_1 - r_2 + s_2 - 1 &= r_2 - d_2 + 1 \end{cases}$$

Using also (4.1) this is the case if and only if the following 3 equations hold

$$r_2 = r_1 + d_2 - d_1 \tag{4.2}$$

$$s_1 = (3 - d_2)r_1 + (d_2 - 3)d_1 + 2 \tag{4.3}$$

$$s_2 = (3 - d_1)r_1 + (d_1 - 3)d_1 + 2 \tag{4.4}$$

Case 1: $d_1 > d_2 \geq 4$.

By (4.4) and since $d_1 \geq 5$ we have $r_1 \leq d_1 + \frac{2}{d_1-3} \leq d_1 + 1$. On the other hand (4.2) implies $r_1 \geq d_1 - d_2 + 1$. Hence we obtain

$$r_1 = d_1 - d_2 + i \text{ with } 1 \leq i \leq d_2 + 1$$

and thus

$$\begin{aligned} \dim P &= d_2 r_1 - d_1 d_2 + d_1 + d_2 - r_1 - r_2 + s_1 - 1 \\ &= i - d_2 + 1 \leq 2 \end{aligned}$$

Hence there is no Prym-Tyurin variety P of dimension ≥ 3 in this case.

Case 2: $d_1 \geq 4, d_2 = 3$.

Here we have by (4.4) $r_1 \leq d_1 + \frac{2}{d_1-3} \leq d_1 + 2$, whereas (4.2) gives $r_1 \geq d_1 - 3$. Hence we get

$$r_1 = d_1 - 3 + i \text{ with } 0 \leq i \leq 5$$

and thus using that $s_1 = 2$ by (4.3),

$$\begin{aligned} \dim P &= d_2 r_1 - d_1 d_2 + d_1 + d_2 - r_1 - r_2 + 1 \\ &= r_1 - d_1 + 1 = i - 2 \leq 3. \end{aligned}$$

Hence in this case there is no Prym-Tyurin variety P of dimension ≥ 4 .

Case 3: $d_1 \geq 4, d_2 = 2$.

Again we have by (4.4) $r_1 \leq d_1 + \frac{2}{d_1-3}$. But now $d_1 \geq 5$ since $(d_1, d_2) = 1$, implying $r_1 \leq d_1 + 1$. As above we get $r_1 \geq d_1 - 3$ and hence

$$r_1 = d_1 - 3 + i \text{ with } 0 \leq i \leq 4$$

and thus using that $s_1 = r_2$ by (4.2) and (4.3)

$$\begin{aligned} \dim P &= r_1 - d_1 + 1 \\ &= i - 2 \leq 2. \end{aligned}$$

Hence in this case there no Prym-Tyurin variety of dimension ≥ 3 .

Case 4: $d_1 = 3, d_2 = 2$.

Here equations (4.2) - (4.4) say $s_1 = r_2 = r_1 - 1$ and $s_2 = 2$ and thus are fulfilled for any $r_1 \geq 2$. According to what we have said above P is a Prym-Tyurin variety of exponent 6 in J_X . Its dimension is $\dim P = r_1 - 2$. $\square$

Remark 4.2. *In Theorem 4.1 we obtain for every dimension $d \geq 4$ a family of Prym-Tyurin varieties of dimension d and exponent 6. Counting parameters it is easy to see that the family is of dimension $2d + 1$.*

Theorem 4.3. *Let the assumptions be as in Proposition 3.4. In particular we assume that $\gcd(d_1, d_2) = 1$, that f_1 does not factorize via a cyclic étale covering and that f_2 is étale cyclic.*

Then $P = P(f_1, f_2)$ is a Prym-Tyurin variety of dimension ≥ 5 in J_X if and only if

$$d_1 = 3, d_2 = 2, r_1 = r_2 = s_1 \geq 7 \text{ and } s_2 = 0.$$

In this case P is a Prym-Tyurin variety of dimension $r_1 - 2$ and of exponent 6 in J_X .

Proof. According to Proposition 3.4 P is a Prym-Tyurin variety of exponent $d_1 d_2$ in J_X if and only if $\dim P = g(X_1) = g(X_2) - 1$ and $d_1 = 3$. Using (4.1) this is the case if and only if

$$\begin{cases} d_1 = 3 \text{ and } s_2 = 0 \\ r_2 = d_2 + r_1 - 2 \\ s_1 = 3r_2 - d_2 r_1 \end{cases} \quad (4.5)$$

Now $s_1 \geq 0$ and hence (4.5) implies

$$(d_2 - 3)(3 - r_1) + 3 = 3d_2 + 3r_1 - 6 - d_2 r_1 \geq 0$$

On the other hand $r_1 > 0$ and $r_2 > 0$, since there are no nontrivial unramified coverings of $\mathbb{P}^1$.

If $d_2 \geq 7$, this gives $r_1 \leq 3$ and thus $\dim P = r_1 - 2 \leq 1$. If $d_2 = 6$ or 5, then $r_1 \leq 4$ and thus $\dim P = r_1 - 2 \leq 2$. If $d_2 = 4$, then $r_1 \leq 6$ and thus $\dim P = r_1 - 2 \leq 4$. Hence in these cases there is no Prym-Tyurin

variety P of dimension ≥ 5 .

We are left with $d_2 = 2$ which leads to the conditions of the theorem. According to Proposition 3.4 in this case P is a Prym-Tyurin variety of dimension $r_1 - 2$. $\square$

Remark 4.4. *It is easy to see that in the case of Theorem 4.3 the composed covering $g_i \circ f_i : X \rightarrow \mathbb{P}^1$ is Galois with Galois group S_3 . Again we obtain for every dimension $d \geq 5$ a family of Prym-Tyurin varieties of dimension d and exponent 6. Counting parameters it is easy to see that the family is of dimension $2d + 1$.*

Remark 4.5. *It is easy to see that these are the only cases of pairs of morphisms (f_1, f_2) as in diagram (3.1) where we get a Prym-Tyurin variety in this way.*

5. THE ABEL-PRYM MAP

Let the notation be as in Theorems 4.1 or 4.3. In particular $P = P(f_1, f_2)$ is the Prym-Tyurin variety of the pair (f_1, f_2) . Let $\alpha : X \rightarrow J_X$ denote the Abel map with respect to a base point x_0 . Identifying J_X and P with their dual abelian varieties via the principal polarizations, the canonical embedding $\iota_P : P \rightarrow J_X$ induces a surjective homomorphism $\hat{\iota} : J_X \rightarrow P$. The composed map

$$\beta : X \xrightarrow{\alpha} J_X \xrightarrow{\hat{\iota}} P$$

is called the *Abel-Prym map* of the Prym-Tyurin variety P . Let $R_1 \subset X$ denote the branch locus of the double covering $f_1 : X \rightarrow X_1$.

Proposition 5.1. *If $\dim P \geq 8$, then we have for the Abel-Prym map $\beta : X \rightarrow P$:*

β is injective on $X \setminus R_1$ and contracts R_1 to a point in $P \setminus \beta(X \setminus R_1)$.

Proof. The Prym-Tyurin variety P is the complement of the abelian subvariety $f_1^* J_{X_1} + f_2^* J_{X_2}$ of J_X with respect to the canonical polarization. $f_1^* J_{X_1} + f_2^* J_{X_2}$ is of exponent 6 in J_X . Let $N_{f_i} : J_X \rightarrow J_{X_i}$ denote the Norm map for $i = 1, 2$. It is easy to see that

$$\hat{\iota} = 6 \cdot 1_{J_X} - 3N_{f_1} - 2N_{f_2}$$

Assume that there are points $x \neq y$ of X such that $\beta(x) = \beta(y)$. If ι_1 denotes the involution of X induced by f_1 and for any $x \in X$ we denote by $\iota_2(x)$ and $\iota_2^2(x)$ the other 2 points of the fibre $f_2^{-1}(f_2(x))$ (note that ι_2 is not a global automorphism of X , but locally this notation makes sense), this means

$$6x - 3(1 + \iota_1)(x) - 2(1 + \iota_2 - \iota_2^2)(x) \sim 6y - 3(1 + \iota_1)(y) - 2(1 + \iota_2 - \iota_2^2)(y)$$

or equivalently

$$x + 3\iota_1(y) + 2\iota_2(y) + 2\iota_2^2(y) \sim y + 3\iota_1(x) + 2\iota_2(x) + 2\iota_2^2(x) \quad (6.1)$$

Here $\sim$ means linear equivalence of divisors on the curve X .

It suffices to show that if x or $y \notin R_1$ this implies that (6.1) defines a pencil of divisors of degree 8, since then

either the g_8^1 does not factor via f_1 in which case Castelnuovo's formula gives $g(X) \leq 1 \cdot 7 + 2g(X_1)$ giving $3r_1 - 6 \leq 7 + 2r_1 - 4$ which implies $r_1 \leq 9$,

or the g_8^1 factors via f_1 in which case Castelnuovo's formula applied to X_1 gives $r_1 - 2 = g(X_1) \leq 2 \cdot 3$ which implies $r_1 \leq 8$. But by assumption $r_1 = \dim P + 2 \geq 10$, a contradiction.

In order to complete the proof, assume (6.1) is an equality of divisors.

Assume first that x and y are in the same fibre of the composed map $g_i \circ f_i : X \rightarrow \mathbb{P}^1$. Since certainly $3\iota_1(y) \neq 3\iota_1(x)$ we may assume after eventually exchanging $\iota_2(x)$ and $\iota_2^2(x)$ respectively $\iota_2(y)$ and $\iota_2^2(y)$ that

$$3\iota_1(y) = y + 2\iota_2(x) \quad \text{and} \quad 3\iota_1(x) = x + 2\iota_2(y)$$

In particular x and y are two different ramification points of f_1 in a fibre of $g_1 \circ f_1$. But every fibre of $g_1 \circ f_1$ contains at most one ramification point of f_1 , a contradiction.

If x and y are not in the same fibre of $g_i \circ f_i$ and f_1 is not ramified at x or y , then (6.1) cannot be an equality of divisors and the same argument works. So finally assume that x and y are different points of R_1 . Then (6.1) reads

$$2y + 2\iota_2(y) + 2\iota_2^2(y) \sim 2x + 2\iota_2(x) + 2\iota_2^2(x)$$

which means that (6.1) is the g_6^1 defined by $g_i \circ f_i$. In particular $\beta(R_1)$ contracts to a point. $\square$

Remark 5.2. *Observe that with the notation of the proof of Proposition 5.1 the correspondence D_1 of the curve C defined by*

$$D_1(x) = x - 3\iota_1(x) - 2\iota_2(x) - 2\iota_2^2(x)$$

for any $x \in X$ induces the endomorphism $\hat{i}$ with image P . The correspondence D of C defined by

$$D(x) = \pi^*\pi(x) - D_1(x)$$

is an effective correspondence defining $\hat{i}$. But D is not fixed point free, so one cannot apply Kanev's theorem (see [1], Theorem 12.9.1) in order to show that P is a Prym-Tyurin variety.

Remark 5.3. *One can show that the singularity $\beta(R_1)$ of the curve $\beta(X)$ is an ordinary singularity. We do not include the details here.*

Remark 5.4. *Recall that the Seshadri constant of the polarisation Ξ of the abelian variety P is defined as*

$$\varepsilon(\Xi) = \inf \frac{(\Xi \cdot C)}{\text{mult}_0(C)}$$

where the multiplicity is to be taken over all irreducible reduced curves C on P containing 0 . The existence of a curve with high multiplicity means that the Seshadri constant is small.

In fact

$$\varepsilon(\Xi) \leq \begin{cases} 3 - \frac{5}{\dim P + 1} & \text{if } P \text{ is as in Theorem 4.1,} \\ 3 - \frac{5}{\dim P + 2} & \text{Theorem 4.3.} \end{cases}$$

This follows from the fact that C is of class $\frac{6}{(\dim P - 1)!} [\Xi]^{\dim P - 1}$.

REFERENCES

- [1] Birkenhake, Ch., Lange, H.: Complex Abelian Varieties. Springer, Grundlehren 302, 2nd edition (2004).
- [2] Lange, H. Pauly, C.: Abelianization of moduli spaces of principal bundles (to appear).
- [3] Lange, H. Recillas, S.: Prym varieties of pairs of coverings, Adv. in Geometry (2004)

H. LANGE, MATHEMATISCHES INSTITUT, UNIVERSITÄT ERLANGEN-NÜRNBERG,
BISMARCKSTRASSE 1 $\frac{1}{2}$, D-91054 ERLANGEN, GERMANY
E-mail address: lange@mi.uni-erlangen.de